

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Sinskey *et al.*

Appl. No. To be assigned
(Divisional of U.S. Appl. No. 09/677,575;
Filed: October 3, 2000)

Filed: Herewith

For: **Pyruvate Carboxylase from
*Corynebacterium glutamicum***

Art Unit: To be Assigned

Examiner: To be Assigned

Atty. Docket: 1533.0790002/MAC/AGU

Preliminary Amendment and Submission of Sequence Listing

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Prior to examination on the merits, kindly amend the captioned application as follows.

This Preliminary Amendment is provided in the following format:

(A) A clean version of each replacement paragraph/section/claim along with clear instructions for entry;

(B) Starting on a separate page, appropriate remarks and arguments. *See* 37 C.F.R. § 1.121 and MPEP 714; and

(C) Starting on a separate page, a marked-up version entitled: "Version with markings to show changes made."

It is not believed that extensions of time or fees for net addition of claims are required beyond those that may otherwise be provided for in documents accompanying this paper. However, if additional extensions of time are necessary to prevent abandonment of this application, then such extensions of time are hereby petitioned under 37 C.F.R. § 1.136(a), and any fees required therefor (including

fees for net addition of claims) are hereby authorized to be charged to our Deposit Account No. 19-0036.

Amendments

In the Specification:

Please insert the sequence listing at the end of the application.

Please substitute the paragraph beginning on page 2, line 21, with the following paragraph:

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a pyruvate carboxylase polypeptide having the amino acid sequence in Figures 1A-C (SEQ ID NO:2) or the amino acid sequence encoded by the clone deposited in a bacterial host as ATCC Deposit Number PTA-982. The nucleotide sequence determined by sequencing the deposited pyruvate carboxylase clone, which is shown in Figures 1A-C (SEQ ID NO:1), contains an open reading frame encoding a polypeptide of 1140 amino acid residues which has a deduced molecular weight of about 123.6 kDa. The 1140 amino acid sequence of the predicted pyruvate carboxylase protein is shown in Figures 1A-C and in SEQ ID NO:2.

Please substitute the paragraph beginning on page 2, line 31, with the following paragraph:

Thus, one aspect of the invention provides an isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence in SEQ ID NO:2; (b) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the clone contained in ATCC Deposit No. PTA-982; and (c) a nucleotide sequence complementary to any of the nucleotide sequences in (a) or (b) above.

Please substitute the paragraph beginning on page 3, line 21, with the following paragraph:

The invention further provides an isolated pyruvate carboxylase polypeptide having amino acid sequence selected from the group consisting of: (a) the amino acid sequence of the pyruvate carboxylase polypeptide having the amino acid sequence shown in Figures 1A-C (SEQ ID NO:2); and (b) the amino acid sequence of the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the clone contained in ATCC Deposit No. PTA-982. The polypeptides of the present invention also include polypeptides having an amino acid sequence with at least 90% similarity, more preferably at least 95% similarity to those described in (a) or (b) above, as well as polypeptides having an amino acid sequence at least 70% identical, more preferably at least 90% identical, and still more preferably 95%, 97%, 98% or 99% identical to those above.

Please substitute the paragraph beginning on page 4, line 3, with the following paragraph:

Figures 1A-C show the nucleotide (SEQ ID NO:1) and deduced amino acid (SEQ ID NO:2) sequences of the complete pyruvate carboxylase protein determined by sequencing of the DNA clone contained in ATCC Deposit No. PTA-982. The protein has sequence of about 1140 amino acid residues and a deduced molecular weight of about 123.6 kDa.

Please substitute the paragraph beginning on page 4, line 11, with the following paragraph:

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding the pyruvate carboxylase protein having the amino acid sequence shown in Figures 1A-C (SEQ ID NO:2) which was determined by sequencing a cloned cosmid. The pyruvate carboxylase protein of the present invention shares sequence homology with *M. tuberculosis* and human pyruvate carboxylase proteins. The nucleotide sequence shown in Figures 1A-C (SEQ ID NO:1) was obtained by sequencing cosmid III F10 encoding a pyruvate carboxylase polypeptide. A clone containing the pyruvate carboxylase gene was deposited on November 22, 1999 at the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, and given accession number PTA-982.

Please substitute the paragraph beginning on page 5, line 22, with the following paragraph:

Using the information provided herein, such as the nucleotide sequence in Figures 1A-C, a nucleic acid molecule of the present invention encoding a pyruvate carboxylase polypeptide may be obtained using standard cloning and screening procedures, such as those for cloning DNAs using mRNA as starting material. The pyruvate carboxylase protein shown in Figures 1A-C (SEQ ID NO:2) is about 63% identical to *M. tuberculosis* and 44% identical to human. As one of ordinary skill would appreciate, due to the possibilities of sequencing errors discussed above, as well as the variability of cleavage sites for leaders in different known proteins, the actual pyruvate carboxylase polypeptide encoded by the deposited clone comprises about 1140 amino acids, but may be anywhere in the range of 1133-1147 amino acids.

Please substitute the paragraph beginning on page 6, line 16, with the following paragraph:

Isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) with an initiation codon at positions 199-201 of the nucleotide sequence shown in Figures 1A-C (SEQ ID NO:1); DNA molecules comprising the coding sequence for the pyruvate carboxylase protein shown in Figures 1A-C and SEQ ID NO:2; and DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the pyruvate carboxylase protein. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate the degenerate variants described above.

Please substitute the paragraph beginning on page 6, line 25, with the following paragraph:

In another aspect, the invention provides isolated nucleic acid molecules encoding the pyruvate carboxylase polypeptide having an amino acid sequence encoded by the clone deposited as ATCC Deposit No. PTA-982. Preferably, this nucleic acid molecule will encode the polypeptide encoded by the above-described deposited clone. The invention further provides an isolated nucleic acid molecule having the nucleotide sequence shown in Figures 1A-C (SEQ ID NO:1) or the

nucleotide sequence of the pyruvate carboxylase DNA contained in the above-described deposited clone, or nucleic acid molecule having a sequence complementary to one of the above sequences.

Please substitute the paragraph beginning on page 7, line 2, with the following paragraph:

In another aspect, the invention provides an isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above, for instance, the clone contained in ATCC Deposit PTA-982. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150 mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C. By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers.

Please substitute the paragraph beginning on page 7, line 16, with the following paragraph:

Of course, polynucleotides hybridizing to a larger portion of the reference polynucleotide (e.g., the deposited clone), for instance, a portion 50-750 nt in length, or even to the entire length of the reference polynucleotide, also useful as probes according to the present invention, as are polynucleotides corresponding to most, if not all, of the nucleotide sequence of the deposited DNA or the nucleotide sequence as shown in Figures 1A-C (SEQ ID NO:1). By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide, (e.g., the deposited DNA or the nucleotide sequence as shown in Figures 1A-C (SEQ ID NO:1)). As indicated, such portions are useful diagnostically either as a probe according to conventional DNA hybridization techniques or as primers for amplification of a target sequence by the polymerase chain reaction (PCR), as described, for instance, in *Molecular Cloning, A Laboratory Manual*, 2nd. edition, edited by

Sambrook, J., Fritsch, E. F. and Maniatis, T., (1989), Cold Spring Harbor Laboratory Press, the entire disclosure of which is hereby incorporated herein by reference.

Please substitute the paragraph beginning on page 7, line 32, with the following paragraph:

Since a pyruvate carboxylase clone has been deposited and its determined nucleotide sequence is provided in Figures 1A-C (SEQ ID NO:1), generating polynucleotides which hybridize to a portion of the pyruvate carboxylase DNA molecule would be routine to the skilled artisan. For example, restriction endonuclease cleavage or shearing by sonication of the pyruvate carboxylase clone could easily be used to generate DNA portions of various sizes which are polynucleotides that hybridize to a portion of the pyruvate carboxylase DNA molecule. Alternatively, the hybridizing polynucleotides of the present invention could be generated synthetically according to known techniques.

Please substitute the paragraph beginning on page 9, line 6, with the following paragraph:

Such variants include those produced by nucleotide substitutions, deletions or additions. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the pyruvate carboxylase protein or portions thereof. Also especially preferred in this regard are conservative substitutions. Most highly preferred are nucleic acid molecules encoding the pyruvate carboxylase protein having the amino acid sequence shown in Figures 1A-C (SEQ ID NO:2).

Please substitute the paragraph beginning on page 9, line 19, with the following paragraph:

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 90% identical, and more preferably at least 95%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence in SEQ ID NO:2; (b) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded

by the clone contained in ATCC Deposit No. PTA-982; or (c) a nucleotide sequence complementary to any of the nucleotide sequences in (a) or (b).

Please substitute the paragraph beginning on page 10, line 11, with the following paragraph:

As a practical matter, whether any particular nucleic acid molecule is at least 90%, 95%, 97%, 98% or 99% identical to, for instance, the nucleotide sequence shown in Figures 1A-C or to the nucleotides sequence of the deposited clone can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman (*Advances in Applied Mathematics* 2: 482-489 (1981)) to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

Please substitute the paragraph beginning on page 10, line 25, with the following paragraph:

The present application is directed to nucleic acid molecules at least 90%, 95%, 97%, 98% or 99% identical to the nucleic acid sequence shown in Figures 1A-C (SEQ ID NO:1) or to the nucleic acid sequence of the deposited DNA, irrespective of whether they encode a polypeptide having pyruvate carboxylase activity. This is because, even where a particular nucleic acid molecule does not encode a polypeptide having pyruvate carboxylase activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer.

Please substitute the paragraph beginning on page 11, line 1, with the following paragraph:

Preferred, however, are nucleic acid molecules having sequences at least 90%, 95%, 97%, 98% or 99% identical to the nucleic acid sequence shown in Figures 1A-C (SEQ ID NO:1) or to the

nucleic acid sequence of the deposited DNA which do, in fact, encode a polypeptide having pyruvate carboxylase protein activity. By "a polypeptide having pyruvate carboxylase activity" is intended polypeptides exhibiting activity similar, but not necessarily identical, to an activity of the pyruvate carboxylase protein of the invention as measured in a particular biological assay.

Please substitute the paragraph beginning on page 11, line 9, with the following paragraph:

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 90%, 95%, 97%, 98%, or 99% identical to the nucleic acid sequence of the deposited DNA or the nucleic acid sequence shown in Figures 1A-C (SEQ ID NO:1) will encode a polypeptide "having pyruvate carboxylase protein activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having pyruvate carboxylase protein activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid).

Please substitute the paragraph beginning on page 15, line 15, with the following paragraph:

The invention further provides an isolated pyruvate carboxylase polypeptide having the amino acid sequence encoded by the deposited DNA, or the amino acid sequence in Figures 1A-C (SEQ ID NO:2), or a peptide or polypeptide comprising a portion of the above polypeptides. The terms "peptide" and "oligopeptide" are considered synonymous (as is commonly recognized) and each term can be used interchangeably as the context requires to indicate a chain of at least to amino acids coupled by peptidyl linkages. The word "polypeptide" is used herein for chains containing more than ten amino acid residues. All oligopeptide and polypeptide formulas or sequences herein are written from left to right and in the direction from amino terminus to carboxy terminus.

Please substitute the paragraph beginning on page 17, line 22, with the following paragraph:

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 97%, 98% or 99% identical to, for instance, the amino acid sequence shown in Figures 1A-C (SEQ ID NO:2) or to the amino acid sequence encoded by deposited clone can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

Please substitute the paragraph beginning on page 24, line 29, with the following paragraph:

PCR was performed using the Boehringer Mannheim PCR core kit following the manufacturer's instructions. When PCR was performed on *Corynebacterium* chromosomal DNA, about 1 μ g DNA was used in each reaction. The forward primer used was 5'GTCTTCATCGAGATGAATCCGCG3' (SEQ ID NO:3) and the reverse primer used was 5'CGCAGCGCCACATCGTAAGTCGC3' (SEQ ID NO:4) for the PCR reaction.

Please substitute table 1 beginning on page 27, line 16, with the following table 1:

Organism	Conserved region A	Conserved region B
<i>Caenorhabditis elegans</i>	YFIEVNAR (SEQ ID NO:5)	ATFDVSM (SEQ ID NO:6)
<i>Aedes aegypti</i>	YFIEVNAR (SEQ ID NO:7)	ATFDVAL (SEQ ID NO:8)
<i>Mycobacterium tuberculosis</i>	VFIEMNPR (SEQ ID NO:9)	ATYDVAL (SEQ ID NO:10)
<i>Bacillus stearothermophilus</i>	YFIEVNPR (SEQ ID NO:11)	ATFDVAY (SEQ ID NO:12)
<i>Pichia pastoris</i>	YFIEINPR (SEQ ID NO:13)	ATFDVSM (SEQ ID NO:14)
<i>Mus musculus</i>	YFIEVNRSR (SEQ ID NO:15)	ATFDVAM (SEQ ID NO:16)
<i>Rattus norvegicus</i>	YFIEVNRSR (SEQ ID NO:17)	ATFDVAM (SEQ ID NO:18)
<i>Saccharomyces cerevisiae 1</i>	YFIEINPR (SEQ ID NO:19)	ATFDVAM (SEQ ID NO:20)
<i>Saccharomyces cerevisiae 2</i>	YFIEINPR (SEQ ID NO:21)	ATFDVAM (SEQ ID NO:22)
<i>Rhizabium etli</i>	YFIEVNPR (SEQ ID NO:23)	ATFDVSM (SEQ ID NO:24)
<i>Homo sapiens</i>	YFIEVNRSR (SEQ ID NO:25)	ATFDVAM (SEQ ID NO:26)
<i>Schizosaccharomyces pombe</i>	YFIEINPR (SEQ ID NO:27)	ATFDVSM (SEQ ID NO:28)

Please substitute the paragraph beginning on page 28, line 25, with the following paragraph:

The amino-terminal segment of the *C. glutamicum* pyruvate carboxylase contains the hexapeptide GGGGRG (SEQ ID NO:37), which matches the GGGG(R/K)G (SEQ ID NO:38) sequence that is found in all biotin-binding proteins and is believed to be an ATP-binding site (Fry, D.C., *et al.*, *Proc. Natl. Acad. Sci. USA* 83:907-911 (1986); Post, L.E., *et al.*, *J. Biol. Chem.* 265:7742-7747 (1990)). A second region that is proposed to be involved in ATP binding and is present in biotin-dependent carboxylases and carbamylphosphate synthetase (Lim, F., *et al.*, *J. Biol. Chem.* 263:11493-11497 (1988)) is conserved in the *C. glutamicum* sequence. The predicted *C. glutamicum* pyruvate carboxylase protein also contains a putative pyruvate-binding motif, FLFEDPWDR (SEQ ID NO:29), which is conserved in the transcarboxylase domains of *Mycobacterium*, *Rhizobium* and human pyruvate carboxylases (Dunn, M.F., *et al.*, *J. Bacteriol.* 178:5960-5970 (1996)). Tryptophan fluorescence studies with transcarboxylase have shown that the Trp residue present in this motif is involved in pyruvate binding (Kumer, G.K., *et al.*, *Biochemistry* 27:5978-5983 (1988)). The carboxy-terminal segment of the enzyme contains a putative biotin-binding site, AMKM (SEQ ID NO:39), which is identical to those found in other pyruvate carboxylases as well as the biotin-carboxyl-carrier protein (BCCP) domains of other biotin-dependent enzymes.

Please substitute table 2 beginning on page 29, line 9, with the following table 2:

<i>Primer name</i>	<i>Primer sequence (5'-3')</i>
<i>Begrev1</i>	<i>TTCACCAGGTCCACCTCG</i> (SEQ ID NO:30)
<i>Endfor1</i>	<i>CGTCGCAAAGCTGACTCC</i> (SEQ ID NO:31)
<i>Begrev2</i>	<i>GATGCTTCTGTTGCTAATTTGC</i> (SEQ ID NO:32)
<i>Endfor2</i>	<i>GGCCATTAAGGATATGGCTG</i> (SEQ ID NO:33)
<i>Begrev3</i>	<i>GCGGTGGAATGATCCCCGA</i> (SEQ ID NO:34)
<i>Endfor3</i>	<i>ACCGCACTGGGCCTTGCG</i> (SEQ ID NO:35)
<i>Endfor4</i>	<i>TCGCCGCTTCGGCAACAC</i> (SEQ ID NO:36)

In the Claims:

Please cancel claims 1-13 without prejudice to or disclaimer of the subject matter encompassed thereby.

Please amend the claims as follows:

14. (Once Amended) An isolated pyruvate carboxylase polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

(a) the amino acid sequence of the pyruvate carboxylase polypeptide having the complete amino acid sequence in SEQ ID NO:2; and

(b) the amino acid sequence of the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the clone contained in ATCC Deposit No. PTA-982.

15. (Once Amended) A method of making amino acids comprising expressing an isolated nucleic acid molecule encoding pyruvate carboxylase comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the amino acid sequence in SEQ ID NO:2;
- (b) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the clone contained in ATCC Deposit No. PTA-982; and
- (c) a nucleotide sequence complementary to any of the nucleotide sequences in (a) or (b), and recovering said amino acids.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Remarks

In the documents accompanying this Preliminary Amendment, claims 1-13 have been canceled. Upon entry of the foregoing amendment, claims 14-17 are pending in the application, with claims 14 and 15 being the independent claims. These changes are believed to introduce no new matter, and their entry is respectfully requested.

Applicants have amended the specification to refer to the ATCC Deposit No. of a deposited plasmid clone containing the pyruvate carboxylase gene and to direct the entry of the formal drawings where appropriate. Applicants have amended the specification to delete reference to a deposited "cosmid" clone. The specification has also been amended to direct the entry of this sequence listing after the claims of the above identified application and to provide the SEQ ID NO's next to the specific sequence. These changes are believed to introduce no new matter, and their entry is respectfully requested.

Claim 14 has been amended to correct a typographical error and is believed to introduce no new matter. Claim 15 has been amended to delete reference to a canceled claim, and to insert the contents of the canceled claim.

In accordance with 37 C.F.R. § 1.821(g), this submission includes no new matter.

In accordance with 37 C.F.R. § 1.821(f), the paper copy of the Sequence Listing and the computer readable copy of the Sequence Listing submitted herewith in the above application are the same.

Conclusion

Applicants believe the present application is in condition for allowance. If the Examiner believes, for any reason, that personal communication will expedite prosecution of this application, the Examiner is invited to telephone the undersigned at the number provided.

Prompt and favorable consideration of this Preliminary Amendment is respectfully requested.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Michele A. Cimbala
Attorney for Applicants
Registration No. 33,851

Date: January 15, 2002

1100 New York Avenue, N.W.
Suite 600
Washington, D.C. 20005-3934
(202) 371-2600

Version with markings to show changes made

In the Specification:

Please substitute the paragraph beginning on page 2, line 21, with the following paragraph:

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a pyruvate carboxylase polypeptide having the amino acid sequence in Figures 1A-C (SEQ ID NO:2) or the amino acid sequence encoded by the [cosmid] clone deposited in a bacterial host as ATCC Deposit Number [____] PTA-982. The nucleotide sequence determined by sequencing the deposited pyruvate carboxylase [cosmid] clone, which is shown in Figures 1A-C (SEQ ID NO:1), contains an open reading frame encoding a polypeptide of 1140 amino acid residues which has a deduced molecular weight of about 123.6 kDa. The 1140 amino acid sequence of the predicted pyruvate carboxylase protein is shown in Figures 1A-C and in SEQ ID NO:2.

Please substitute the paragraph beginning on page 2, line 31, with the following paragraph:

Thus, one aspect of the invention provides an isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence in SEQ ID NO:2; (b) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the [cosmid] clone contained in ATCC Deposit No. [____] PTA-982; and (c) a nucleotide sequence complementary to any of the nucleotide sequences in (a) or (b) above.

Please substitute the paragraph beginning on page 3, line 21, with the following paragraph:

The invention further provides an isolated pyruvate carboxylase polypeptide having amino acid sequence selected from the group consisting of: (a) the amino acid sequence of the pyruvate carboxylase polypeptide having the amino acid sequence shown in Figures 1A-C (SEQ ID NO:2); and (b) the amino acid sequence of the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the [cosmid] clone contained in ATCC Deposit No. [____] PTA-982. The polypeptides of the present invention also include polypeptides having an amino acid

sequence with at least 90% similarity, more preferably at least 95% similarity to those described in (a) or (b) above, as well as polypeptides having an amino acid sequence at least 70% identical, more preferably at least 90% identical, and still more preferably 95%, 97%, 98% or 99% identical to those above.

Please substitute the paragraph beginning on page 4, line 3, with the following paragraph:

Figures 1A-C show[s] the nucleotide (SEQ ID NO:1) and deduced amino acid (SEQ ID NO:2) sequences of the complete pyruvate carboxylase protein determined by sequencing of the DNA clone contained in ATCC Deposit No. [_____] PTA-982. The protein has sequence of about 1140 amino acid residues and a deduced molecular weight of about 123.6 kDa.

Please substitute the paragraph beginning on page 4, line 11, with the following paragraph:

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding the pyruvate carboxylase protein having the amino acid sequence shown in Figures 1A-C (SEQ ID NO:2) which was determined by sequencing a cloned cosmid. The pyruvate carboxylase protein of the present invention shares sequence homology with *M. tuberculosis* and human pyruvate carboxylase proteins. The nucleotide sequence shown in Figures 1A-C (SEQ ID NO:1) was obtained by sequencing cosmid III F10 encoding a pyruvate carboxylase polypeptide, [, which] A clone containing the pyruvate carboxylate gene was deposited on [_____] November 22, 1999 at the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, and given accession number [_____] PTA-982.

Please substitute the paragraph beginning on page 5, line 22, with the following paragraph:

Using the information provided herein, such as the nucleotide sequence in Figures 1A-C, a nucleic acid molecule of the present invention encoding a pyruvate carboxylase polypeptide may be obtained using standard cloning and screening procedures, such as those for cloning DNAs using mRNA as starting material. The pyruvate carboxylase protein shown in Figures 1A-C (SEQ ID NO:2) is about 63% identical to *M. tuberculosis* and 44% identical to human. As one of ordinary skill would appreciate, due to the possibilities of sequencing errors discussed above, as well as the

variability of cleavage sites for leaders in different known proteins, the actual pyruvate carboxylase polypeptide encoded by the deposited [cosmid] clone comprises about 1140 amino acids, but may be anywhere in the range of 1133-1147 amino acids.

Please substitute the paragraph beginning on page 6, line 16, with the following paragraph:

Isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) with an initiation codon at positions 199-201 of the nucleotide sequence shown in Figures 1A-C (SEQ ID NO:1); DNA molecules comprising the coding sequence for the pyruvate carboxylase protein shown in Figures 1A-C and SEQ ID NO:2; and DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the pyruvate carboxylase protein. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate the degenerate variants described above.

Please substitute the paragraph beginning on page 6, line 25, with the following paragraph:

In another aspect, the invention provides isolated nucleic acid molecules encoding the pyruvate carboxylase polypeptide having an amino acid sequence encoded by the [cosmid] clone deposited as ATCC Deposit No. [_____] PTA-982. Preferably, this nucleic acid molecule will encode the polypeptide encoded by the above-described deposited clone. The invention further provides an isolated nucleic acid molecule having the nucleotide sequence shown in Figures 1A-C (SEQ ID NO:1) or the nucleotide sequence of the pyruvate carboxylase DNA contained in the above-described deposited clone, or nucleic acid molecule having a sequence complementary to one of the above sequences.

Please substitute the paragraph beginning on page 7, line 2, with the following paragraph:

In another aspect, the invention provides an isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above, for instance, the [cosmid] clone contained in ATCC Deposit [_____] PTA-982. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x

SSC (150 mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 μ g/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C. By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers.

Please substitute the paragraph beginning on page 7, line 16, with the following paragraph:

Of course, polynucleotides hybridizing to a larger portion of the reference polynucleotide (e.g., the deposited [cosmid] clone), for instance, a portion 50-750 nt in length, or even to the entire length of the reference polynucleotide, also useful as probes according to the present invention, as are polynucleotides corresponding to most, if not all, of the nucleotide sequence of the deposited DNA or the nucleotide sequence as shown in Figures 1A-C (SEQ ID NO:1). By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide, (e.g., the deposited DNA or the nucleotide sequence as shown in Figures 1A-C (SEQ ID NO:1)). As indicated, such portions are useful diagnostically either as a probe according to conventional DNA hybridization techniques or as primers for amplification of a target sequence by the polymerase chain reaction (PCR), as described, for instance, in *Molecular Cloning, A Laboratory Manual*, 2nd. edition, edited by Sambrook, J., Fritsch, E. F. and Maniatis, T., (1989), Cold Spring Harbor Laboratory Press, the entire disclosure of which is hereby incorporated herein by reference.

Please substitute the paragraph beginning on page 7, line 32, with the following paragraph:

Since a pyruvate carboxylase [cosmid] clone has been deposited and its determined nucleotide sequence is provided in Figures 1A-C (SEQ ID NO:1), generating polynucleotides which hybridize to a portion of the pyruvate carboxylase DNA molecule would be routine to the skilled artisan. For example, restriction endonuclease cleavage or shearing by sonication of the pyruvate carboxylase [cosmid] clone could easily be used to generate DNA portions of various sizes which

are polynucleotides that hybridize to a portion of the pyruvate carboxylase DNA molecule. Alternatively, the hybridizing polynucleotides of the present invention could be generated synthetically according to known techniques.

Please substitute the paragraph beginning on page 9, line 6, with the following paragraph:

Such variants include those produced by nucleotide substitutions, deletions or additions. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the pyruvate carboxylase protein or portions thereof. Also especially preferred in this regard are conservative substitutions. Most highly preferred are nucleic acid molecules encoding the pyruvate carboxylase protein having the amino acid sequence shown in Figures 1A-C (SEQ ID NO:2).

Please substitute the paragraph beginning on page 9, line 19, with the following paragraph:

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 90% identical, and more preferably at least 95%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence in SEQ ID NO:2; (b) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the [cosmid] clone contained in ATCC Deposit No. [_____] PTA-982; or (c) a nucleotide sequence complementary to any of the nucleotide sequences in (a) or (b).

Please substitute the paragraph beginning on page 10, line 11, with the following paragraph:

As a practical matter, whether any particular nucleic acid molecule is at least 90%, 95%, 97%, 98% or 99% identical to, for instance, the nucleotide sequence shown in Figures 1A-C or to the nucleotides sequence of the deposited [cosmid] clone can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package,

Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman (*Advances in Applied Mathematics* 2: 482-489 (1981)) to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

Please substitute the paragraph beginning on page 10, line 25, with the following paragraph:

The present application is directed to nucleic acid molecules at least 90%, 95%, 97%, 98% or 99% identical to the nucleic acid sequence shown in Figures 1A-C (SEQ ID NO:1) or to the nucleic acid sequence of the deposited DNA, irrespective of whether they encode a polypeptide having pyruvate carboxylase activity. This is because, even where a particular nucleic acid molecule does not encode a polypeptide having pyruvate carboxylase activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer.

Please substitute the paragraph beginning on page 11, line 1, with the following paragraph:

Preferred, however, are nucleic acid molecules having sequences at least 90%, 95%, 97%, 98% or 99% identical to the nucleic acid sequence shown in Figures 1A-C (SEQ ID NO:1) or to the nucleic acid sequence of the deposited DNA which do, in fact, encode a polypeptide having pyruvate carboxylase protein activity. By "a polypeptide having pyruvate carboxylase activity" is intended polypeptides exhibiting activity similar, but not necessarily identical, to an activity of the pyruvate carboxylase protein of the invention as measured in a particular biological assay.

Please substitute the paragraph beginning on page 11, line 9, with the following paragraph:

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least

90%, 95%, 97%, 98%, or 99% identical to the nucleic acid sequence of the deposited DNA or the nucleic acid sequence shown in Figures 1A-C (SEQ ID NO:1) will encode a polypeptide "having pyruvate carboxylase protein activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having pyruvate carboxylase protein activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid).

Please substitute the paragraph beginning on page 15, line 15, with the following paragraph:

The invention further provides an isolated pyruvate carboxylase polypeptide having the amino acid sequence encoded by the deposited DNA, or the amino acid sequence in Figures 1A-C (SEQ ID NO:2), or a peptide or polypeptide comprising a portion of the above polypeptides. The terms "peptide" and "oligopeptide" are considered synonymous (as is commonly recognized) and each term can be used interchangeably as the context requires to indicate a chain of at least to amino acids coupled by peptidyl linkages. The word "polypeptide" is used herein for chains containing more than ten amino acid residues. All oligopeptide and polypeptide formulas or sequences herein are written from left to right and in the direction from amino terminus to carboxy terminus.

Please substitute the paragraph beginning on page 17, line 22, with the following paragraph:

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 97%, 98% or 99% identical to, for instance, the amino acid sequence shown in Figures 1A-C (SEQ ID NO:2) or to the amino acid sequence encoded by deposited [cosmid] clone can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of

identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

Please substitute the paragraph beginning on page 24, line 29, with the following paragraph:

PCR was performed using the Boehringer Mannheim PCR core kit following the manufacturer's instructions. When PCR was performed on *Corynebacterium* chromosomal DNA, about 1 µg DNA was used in each reaction. The forward primer used was

5'GTCTTCATCGAGATGAATCCGCG3' (SEQ ID NO:3) and the reverse primer used was

5'CGCAGCGCCACATCGTAAGTCGC3' (SEQ ID NO:4) for the PCR reaction.

Please substitute table 1 beginning on page 27, line 16, with the following table 1:

Organism	Conserved region A	Conserved region B
<i>Caenorhabditis elegans</i>	YFIEVNAR (SEQ ID NO:5)	ATFDVSM (SEQ ID NO:6)
<i>Aedes aegypti</i>	YFIEVNAR (SEQ ID NO:7)	ATFDVAL (SEQ ID NO:8)
<i>Mycobacterium tuberculosis</i>	VFIEMNPR (SEQ ID NO:9)	ATYDVAL (SEQ ID NO:10)
<i>Bacillus stearothermophilus</i>	YFIEVNPR (SEQ ID NO:11)	ATFDVAY (SEQ ID NO:12)
<i>Pichia pastoris</i>	YFIEINPR (SEQ ID NO:13)	ATFDVSM (SEQ ID NO:14)
<i>Mus musculus</i>	YFIEVNSR (SEQ ID NO:15)	ATFDVAM (SEQ ID NO:16)
<i>Rattus norvegicus</i>	YFIEVNSR (SEQ ID NO:17)	ATFDVAM (SEQ ID NO:18)
<i>Saccharomyces cerevisiae 1</i>	YFIEINPR (SEQ ID NO:19)	ATFDVAM (SEQ ID NO:20)
<i>Saccharomyces cerevisiae 2</i>	YFIEINPR (SEQ ID NO:21)	ATFDVAM (SEQ ID NO:22)
<i>Rhizabium etli</i>	YFIEVNPR (SEQ ID NO:23)	ATFDVSM (SEQ ID NO:24)
<i>Homo sapiens</i>	YFIEVNSR (SEQ ID NO:25)	ATFDVAM (SEQ ID NO:26)
<i>Schizosaccharomyces pombe</i>	YFIEINPR (SEQ ID NO:27)	ATFDVSM (SEQ ID NO:28)

Please substitute the paragraph beginning on page 28, line 25, with the following paragraph:

The amino-terminal segment of the *C. glutamicum* pyruvate carboxylase contains the hexapeptide GGGGRG (SEQ ID NO:37), which matches the GGGG(R/K)G (SEQ ID NO:38) sequence that is found in all biotin-binding proteins and is believed to be an ATP-binding site (Fry, D.C., *et al.*, *Proc. Natl. Acad. Sci. USA* 83:907-911 (1986); Post, L.E., *et al.*, *J. Biol. Chem.* 265:7742-7747 (1990)). A second region that is proposed to be involved in ATP binding and is

present in biotin-dependent carboxylases and carbamylphosphate synthetase (Lim, F., *et al.*, *J. Biol. Chem.* 263:11493-11497 (1988)) is conserved in the *C. glutamicum* sequence. The predicted *C. glutamicum* pyruvate carboxylase protein also contains a putative pyruvate-binding motif, FLFEDPWDR (SEQ ID NO:29), which is conserved in the transcarboxylase domains of *Mycobacterium*, *Rhizobium* and human pyruvate carboxylases (Dunn, M.F., *et al.*, *J. Bacteriol.* 178:5960-5970 (1996)). Tryptophan fluorescence studies with transcarboxylase have shown that the Trp residue present in this motif is involved in pyruvate binding (Kumer, G.K., *et al.*, *Biochemistry* 27:5978-5983 (1988)). The carboxy-terminal segment of the enzyme contains a putative biotin-binding site, AMKM (SEQ ID NO:39), which is identical to those found in other pyruvate carboxylases as well as the biotin-carboxyl-carrier protein (BCCP) domains of other biotin-dependent enzymes.

Please substitute table 2 beginning on page 29, line 9, with the following table 2:

<i>Primer name</i>	<i>Primer sequence (5'-3')</i>
<i>Begrev1</i>	<i>TTCACCAGGTCCACCTCG</i> (<u>SEQ ID NO:30</u>)
<i>Endfor1</i>	<i>CGTCGCAAAGCTGACTCC</i> (<u>SEQ ID NO:31</u>)
<i>Begrev2</i>	<i>GATGCTTCTGTTGCTAATTTGC</i> (<u>SEQ ID NO:32</u>)
<i>Endfor2</i>	<i>GGCCATTAAGGATATGGCTG</i> (<u>SEQ ID NO:33</u>)
<i>Begrev3</i>	<i>GCGGTGGAATGATCCCCGA</i> (<u>SEQ ID NO:34</u>)
<i>Endfor3</i>	<i>ACCGCACTGGGCCTTGCG</i> (<u>SEQ ID NO:35</u>)
<i>Endfor4</i>	<i>TCGCCGCTTCGGCAACAC</i> (<u>SEQ ID NO:36</u>)

In the Claims:

- (a) Claims 1-13 have been cancelled.
- (b) Claims 14 and 15 are amended as follows:

14. (Once Amended) An isolated pyruvate carboxylase polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

(a) the amino acid sequence of the pyruvate carboxylase polypeptide having the complete amino acid sequence in SEQ ID NO:2; and

(b) the amino acid sequence of the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the [cosmid] clone contained in ATCC Deposit No. [____]; and] PTA-982.

15. (Once Amended) A method of making amino acids comprising expressing [the nucleotide sequence of claim 1] an isolated nucleic acid molecule encoding pyruvate carboxylase comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

(a) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the amino acid sequence in SEQ ID NO:2;

(b) a nucleotide sequence encoding the pyruvate carboxylase polypeptide having the complete amino acid sequence encoded by the clone contained in ATCC Deposit No. PTA-982;
and

(c) a nucleotide sequence complementary to any of the nucleotide sequences in (a) or (b), and recovering said amino acids.